



The use of stable isotopes of hydrogen, carbon and oxygen in oil and carbon in individual fatty acids is presented. A study was performed on olive oil samples from different regions (Brda, Slovenia, Istria, Croatia, and Montenegro). In addition, sensory and chemical analyses were obtained and all the samples were classified in accordance with Regulation EEC/2568/91 and related annexes (the last No.1989/2003) as extra virgin olive oils. All measurements performed on genuine olive oils were used to upgrade the database in 2006, 2007 and 2008.

Materials and methods


- Samples from different varieties including *Istrska belica*, *Leccino*, *Maurino*, *Frantoio* from 2006-2008
- Sensory and chemical analysis (sterols)
- Hydrogen, carbon and oxygen isotope analysis on bulk samples – EA-IRMS, HT/EA pyrolysis IRMS
- Chemical characterization of individual fatty acids: GC-MS
- Carbon isotope analysis of individual fatty acids: GC-C-IRMS



Results




The bulk olive oil samples all have isotopic compositions between -31.6‰ and -29.1‰ and a similar range of $\delta^{13}\text{C}$ values in individual fatty acids. Oils from Istria, Slovenian and Montenegro had slightly lower $\delta^{13}\text{C}$ values and showed greater variability. We also observed an increased variability in fatty acid composition in oils from different cultivars from Montenegro.



In the case of the $\delta^{18}\text{O}$ values, these increased from Brda in Slovenia ($23.2 \pm 0.3\text{‰}$) to Montenegro ($25.4 \pm 0.5\text{‰}$), while $\delta^2\text{H}$ values decreased from $-150 \pm 5\text{‰}$ in Brda to $-153 \pm 3\text{‰}$ Montenegro.

Discussion


Authenticity



The $\delta^{13}\text{C}$ values also depend not only on geographical origin but also on the year of production. In all three sampling periods the ratio of $\delta^{13}\text{C}_{16:0}:\delta^{13}\text{C}_{18:1}$ deviates from 1:1, which unfortunately means that that this ratio cannot be used as an indicator of adulteration that had been anticipated in previous studies.

Acknowledgement

This research was performed within the project "The comparison and development of different methods to determine the authenticity of oil and food products" financially supported by Ministry of Agriculture, Forestry and Food and Slovenian Research Agency.



Determination of geographical origin and authenticity of olive oil using stable isotope approach

Nives Ogrinc¹, Milena Bučar Miklavčič^{2,3}, Grazia Casiello⁴ and Francesco Longobardi⁴

¹Dept. of Environ. Sci., Jožef Stefan Institute, Jamova 39, 1000 Ljubljana, Slovenia

² UP ZRS - University of Primorska, Science and Research Centre of Koper, Garibaldijska 1, 6000 Koper, Slovenia

³ LABS, LLC - Institute for ecology, olive oil and control, Zelena ulica 8, 6310 Izola, Slovenia


⁴Dept. of Chemistry, University di Bari, Via Orabona 4, 70126 Bari, Italy



UNIVERSITÀ
DEGLI STUDI DI BARI
ALDO MORO



Geographical origin



Measurements of isotopic composition of genuine olive oils and fatty acids in combination with $\delta^{18}\text{O}$ and $\delta^2\text{H}$ values were further used to determine the geographical origin of olive oil. The most significant variables were $\delta^{13}\text{C}_{18:1}$, $\delta^{13}\text{C}$ of oil and $\delta^{13}\text{C}_{18:0}$ with the overall variability of 83%.

